

IGF-1 Mouse/Rat ELISA (22-IG1MS-E01)

1. List E, et al. The Role of GH in Adipose Tissue: Lessons from Adipose-Specific GH Receptor Gene-Disrupted Mice. *Molecular Endocrinology*. 2013; 27(3): 524-535.
2. Shohreh R, et al. GH, But Not GHRH, Plays a Role in the Development of Experimental Autoimmune Encephalomyelitis. *Endocrinology*. 2011; 152(10): 3803-3810.
3. Bayod S, et al. Long-term treadmill exercise induces neuroprotective molecular changes in rat brain. *Journal of Applied Physiology*. 2011; 111(5): 1380-1390.
4. Andrade S, et al. Immunization Against Active Ghrelin Using Virus-Like Particles for Obesity Treatment. *Curr Pharm Des*. 2013; 19(36): 6551-6558.
5. Fu X, et al. Intravenous administration of mesenchymal stem cells prevents angiotensin II-induced aortic aneurysm formation in apolipoprotein E-deficient mouse. *J Transl Med*. 2013; 11: 175.
6. Ghosh M, et al. Insulin Like Growth Factor-1 Stimulates Differentiation of ATII Cells to ATI-Like Cells Through Activation of Wnt5a. *Am J Physiol Lung Cell Mol Physiol*. 2013; 305(3): L222-8.
7. Menéndez L, et al. IGF-I increases markers of osteoblastic activity and reduces bone resorption via osteoprotegerin and RANK-ligand. *J Transl Med*. 2013; 11: 271.

IGF-1 Human ELISA (22-IGFHU-E01*)

1. Maliszewski A, et al. Prolonged infusion of amino acids increases leucine oxidation in fetal sheep. *American Physiological Society*. 2012; 302: E1483-E1492.
2. Farmer C, et al. Impact of diet deprivation and subsequent over-allowance during prepuberty. Part 1. Effects on growth performance, metabolite status, and mammary gland development in gilts. *J ANIM SCI*. 2011; 90(3): 863-871.
3. Makimura H, et al. The Association of Growth Hormone Parameters with Skeletal Muscle Phosphocreatine Recovery in Adult Men. *The Journal of Clinical Endocrinology & Metabolism*. 2011; 96(3): 817-823.
4. Sundgren N, et al. Extracellular signal-regulated kinase and phosphoinositol-3 kinase mediate IGF-1 induced proliferation of fetal sheep cardiomyocytes. *American Physiological Society*. 2003; 285: R1481-R1489.
5. Ayyavoo A, et al. Severe Hyperemesis Gravidarum Is Associated With Reduced Insulin Sensitivity in the Offspring in Childhood. *The Journal of Clinical Endocrinology & Metabolism*. 2013; 98(8): 3263-3268.
6. Ayyavoo A, et al. First-born Children Have Reduced Insulin Sensitivity and Higher Daytime Blood Pressure Compared to Later-Born Children. *The Journal of Clinical Endocrinology & Metabolism*. 2013; 98(3): 1248-1253.
7. Chien Y, et al. Myostatin and Insulin-Like Growth Factor I: Potential Therapeutic Biomarkers for Pompe Disease. *PLoS ONE*. 2013; 8(8): e71900.
8. Parfieniuk-Kowarda A, et al. Assessment of serum IGF-1 and adipokines related to metabolic dysfunction in HIV-infected adults. *Cytokine*. 2013; 64(1): 97-112.
9. Ayyavoo A, et al. Pre-Pubertal Children Born Post-Term Have Reduced Insulin Sensitivity and Other Markers of the Metabolic Syndrome. *PLoS One*. 2013; 8(7): e67966.

*Assays are for Research Use Only unless otherwise noted
FDA registered for In Vitro Diagnostic use

IGF-1 Human RIA (22-IGF-R20*)

1. Kerver J, et al. Dietary predictors of the insulin-like growth factor system in adolescent females: results from the Dietary Intervention Study in Children (DISC). *Am J Clin Nutr.* 2010; 91(3): 643-650.
2. Nichols T, et al. Protease-Resistant Insulin-Like Growth Factor (IGF)-Binding Protein-4 Inhibits IGF-I Actions and Neointimal Expansion in a Porcine Model of Neointimal Hyperplasia. *Endocrinology.* 2007; 148(10): 5002-5010.
3. Hansen-Pupp I, et al. Postnatal Decrease in Circulating Insulin-Like Growth Factor-I and Low Brain Volumes in Very Preterm Infants. *The Journal of Clinical Endocrinology & Metabolism.* 2011; 96(4): 1129-1135.
4. Martos-Moreno G, et al. Influence of prematurity and growth restriction on the adipokine profile, IGF1, and ghrelin levels in cord blood: relationship with glucose metabolism. *Eur J Endocrinol.* 2009; 161: 381-389.
5. Hansen-Pupp I, et al. Fresh-Frozen Plasma as a Source of Exogenous Insulin-Like Growth Factor-I in the Extremely Preterm Infant. *The Journal of Clinical Endocrinology & Metabolism.* 2009; 94(2): 477-482.
6. Pratipanawatr T, et al. Effect of IGF-I on FFA and glucose metabolism in control and type 2 diabetic subjects. *American Physiological Society.* 2001; 282: E1360-E1368.
7. Aberg D, et al. Serum IGF-I Levels Correlate to Improvement of Functional Outcome after Ischemic Stroke. *The Journal of Clinical Endocrinology & Metabolism.* 2011; 96(7): E1055-E1064.
8. Lekakou L, et al. Effects of gender and age on hypothalamic-pituitary-adrenal reactivity after pharmacological challenge with low-dose 1- μ g ACTH test: a prospective study in healthy adults. *Clin Endocrinol.* 2013; 79(5): 683-8.
9. Joslowski G, et al. Animal Protein Intakes during Early Life and Adolescence Differ in Their Relation to the Growth Hormone-Insulin-Like-Growth-Factor Axis in Young Adulthood. *J Nutr.* 2013; 143(7): 1147-54.

IGF-2 Human ELISA (22-IG2HU-E01)

1. Ayyavoo A, et al. Severe Hyperemesis Gravidarum Is Associated With Reduced Insulin Sensitivity in the Offspring in Childhood. *The Journal of Clinical Endocrinology & Metabolism.* 2013; 98(8): 3263-3268.
2. Ayyavoo A, et al. First-born Children Have Reduced Insulin Sensitivity and Higher Daytime Blood Pressure Compared to Later-Born Children. *The Journal of Clinical Endocrinology & Metabolism.* 2013; 98(3): 1248-1253.
3. Lou Z, et al. Maternal and Fetal IGF-I and IGF-II Levels, Fetal Growth, and Gestational Diabetes. *The Journal of Clinical Endocrinology & Metabolism.* 2012; 97(5): 1720-1728.
4. Mohlin S, et al. HIF2A and IGF2 Expression Correlates in Human Neuroblastoma Cells and Normal Immature Sympathetic Neuroblasts. *Neoplasia.* 2013; 15(3): 328-34.

IGF-2 Human RIA (22-IGF-R30)

1. Bielohuby M, et al. Influence of pre-analytical conditions on the measurement of components of the GH/IGF axis in rats. *Growth Horm IGF Res.* 2013; 23(5): 141-8.

*Assays are for Research Use Only unless otherwise noted
FDA registered for In Vitro Diagnostic use

IGFBP-1 Human ELISA (22-BP1HU-E01)

1. Ayyavoo A, et al. Severe Hyperemesis Gravidarum Is Associated With Reduced Insulin Sensitivity in the Offspring in Childhood. *The Journal of Clinical Endocrinology & Metabolism*. 2013; 98(8): 3263-3268.
2. Ayyavoo A, et al. First-born Children Have Reduced Insulin Sensitivity and Higher Daytime Blood Pressure Compared to Later-Born Children. *The Journal of Clinical Endocrinology & Metabolism*. 2013; 98(3): 1248-1253.
3. Ayyavoo A, et al. Pre-Pubertal Children Born Post-Term Have Reduced Insulin Sensitivity and Other Markers of the Metabolic Syndrome. *PLoS One*. 2013; 8(7): e67966.
4. Joslowski G, et al. Animal Protein Intakes during Early Life and Adolescence Differ in Their Relation to the Growth Hormone-Insulin-Like-Growth-Factor Axis in Young Adulthood. *J Nutr*. 2013; 143(7): 1147-54.

IGFBP-2 Mouse/Rat ELISA (22-BP2MS-E01)

1. Bielohuby M, et al. Serum IGF-I Is Not a Reliable Pharmacodynamic Marker of Exogenous Growth Hormone Activity in Mice. *Endocrinology*. 2011; 152(12): 4764-4776.
2. Castañeda-Gutiérrez E, et al. Effect of a mixture of bovine milk oligosaccharides, *Lactobacillus rhamnosus* NCC4007 and long-chain polyunsaturated fatty acids on catch-up growth of intra-uterine growth-restricted rats. *Acta Physiol*. 2013; doi: 10.1111/apha.12145.
3. Bielohuby M, et al. Influence of pre-analytical conditions on the measurement of components of the GH/IGF axis in rats. *Growth Horm IGF Res*. 2013; 23(5): 141-8.

IGFBP-2 Human ELISA (22-BP2HU-E01)

1. Ayyavoo A, et al. Severe Hyperemesis Gravidarum Is Associated With Reduced Insulin Sensitivity in the Offspring in Childhood. *The Journal of Clinical Endocrinology & Metabolism*. 2013; 98(8): 3263-3268.
2. Ayyavoo A, et al. First-born Children Have Reduced Insulin Sensitivity and Higher Daytime Blood Pressure Compared to Later-Born Children. *The Journal of Clinical Endocrinology & Metabolism*. 2013; 98(3): 1248-1253.
3. Bredella M, et al. IGFBP-2 is a negative predictor of cold-induced brown fat and bone mineral density in young non-obese women. *Bone*. 2013; 53(2):336-9.
4. Joslowski G, et al. Animal Protein Intakes during Early Life and Adolescence Differ in Their Relation to the Growth Hormone-Insulin-Like-Growth-Factor Axis in Young Adulthood. *J Nutr*. 2013; 143(7): 1147-54

*Assays are for Research Use Only unless otherwise noted
FDA registered for In Vitro Diagnostic use

IGFBP-3 Mouse/Rat ELISA (22-BP3MS-E01)

1. Blumbacha K, et al. Dwarfism in Mice Lacking Collagen-binding Integrins $\alpha 2\beta 1$ and $\alpha 11\beta 1$ is Caused by Severely Diminished IGF-1 Levels. *The Journal of Biological Chemistry*. 2012; 287: 6431-6440.
2. Bielohuby M, et al. Serum IGF-I Is Not a Reliable Pharmacodynamic Marker of Exogenous Growth Hormone Activity in Mice. *Endocrinology*. 2011; 152(12): 4764-4776.
3. Birkenfeld A, et al. Influence of the Hepatic Eukaryotic Initiation Factor 2 α (eIF2 α) Endoplasmic Reticulum (ER) Stress Response Pathway on Insulin-mediated ER Stress and Hepatic and Peripheral Glucose Metabolism. *The Journal of Biological Chemistry*. 2011; 286: 36163-36170.
4. Bielohuby M, et al. Lack of Dietary Carbohydrates Induces Hepatic Growth Hormone (GH) Resistance in Rats. *Endocrinology*. 2011; 152(5): 1948-1960.
5. Bielohuby M, et al. Influence of pre-analytical conditions on the measurement of components of the GH/IGF axis in rats. *Growth Horm IGF Res*. 2013; 23(5): 141-8.
6. Haley S, et al. Tactile and Kinesthetic Stimulation (TKS) intervention improves outcomes in weanling rat bone in a neonatal stress model. *J Musculoskelet Neuronal Interact*. 2013; 13(2): 157-165.

IGFBP-3 Human ELISA (22-BP3HU-E01)

1. Buscher A, et al. Functional and total IGFBP3 for the assessment of disorders of the GH/IGF1 axis in children with chronic kidney disease, GH deficiency, or short stature after SGA status at birth. *Eur J Endocrinol*. 2012; 166: 923-931.
2. Ayyavoo A, et al. Severe Hyperemesis Gravidarum Is Associated With Reduced Insulin Sensitivity in the Offspring in Childhood. *The Journal of Clinical Endocrinology & Metabolism*. 2013; 98(8): 3263-3268.
3. Ayyavoo A, et al. First-born Children Have Reduced Insulin Sensitivity and Higher Daytime Blood Pressure Compared to Later-Born Children. *The Journal of Clinical Endocrinology & Metabolism*. 2013; 98(3): 1248-1253.
4. Sampson N, et al. Stromal insulin-like growth factor binding protein 3 (IGFBP3) is elevated in the diseased human prostate and promotes ex vivo fibroblast-to-myofibroblast differentiation. *Endocrinology*. 2013; 154(8): 2586-99.

IGFBP-3 Human RIA (22-IGF-R10)

1. Joslowski G, et al. Animal Protein Intakes during Early Life and Adolescence Differ in Their Relation to the Growth Hormone-Insulin-Like-Growth-Factor Axis in Young Adulthood. *J Nutr*. 2013; 143(7): 1147-54.

Acid-Labile Subunit Human ELISA (22-ALSHU-E01)

1. Hess O, et al. A novel mutation in IGFALS, c.380T>C (p.L127P), associated with short stature, delayed puberty, osteopenia and hyperinsulinaemia in two siblings: insights into the roles of insulin growth factor-1 (IGF1). *Clin Endocrinol*. 2013; doi: 10.1111/cen.12200.
2. Myers J, et al. Integrated proteomics pipeline yields novel biomarkers for predicting preeclampsia. *Hypertension*. 2013; 61(6): 1281-8.

Version 26DEC13

*Assays are for Research Use Only unless otherwise noted
FDA registered for In Vitro Diagnostic use